CLINICAL REASONING GUIDE

Diagnostic Confidence

It takes longer than people think — and that's not necessarily a bad thing.

It doesn't happen the day you graduate, and it doesn't mean already knowing the diagnosis. It means knowing what matters and changing your formulation when the evidence changes.



1 Early on, you feel like you should know

You assess the patient, hear the symptoms, and immediately wonder:

"Is this bipolar disorder?" *"Is this ADHD?"* *"Is this trauma?"*

"What if I'm missing something?"

Then the doubt creeps in: *"Shouldn't I be more sure by now?"*

Uncertainty doesn't automatically mean you're unprepared. Sometimes it means you understand how much information actually matters.

2 One evaluation is a snapshot

At the first visit, you may not have:

- the full symptom timeline
- collateral information
- previous records
- past medication responses
- a clear substance-use history
- enough longitudinal observation



You can't force clarity from information you don't have yet. And the patient may not know — or remember — all of it either.

3 Symptoms overlap. A lot.

The first symptom you notice may point to more than one diagnosis.



Distractibility ≠ ADHD
Mood swings ≠ bipolar
Hallucinations ≠ schizophrenia
Poor focus ≠ depression

The diagnosis comes from the pattern, timeline, context, severity, impairment, and what you've ruled out.

Don't diagnose the symptom. Diagnose the clinical picture.

4 You don't always need certainty

Sometimes the most appropriate answer is: *"This is my working diagnosis based on what I know today."* A working diagnosis lets you:

TREAT	ASSESS	WATCH	REVISE
what needs attention now	what still needs clarifying	how symptoms evolve	when new info changes things

Provisional ≠ careless. It means your formulation is allowed to evolve as you gather more information.

5 Experience changes the questions

Early on: *"What diagnosis fits these symptoms?"* Stronger reasoning adds: *"What else could explain this?"*

MEDICAL	MEDS	SUBSTANCES	OTHER PSYCH	CONTEXT
A medical condition?	Causing or worsening it?	Intoxication, withdrawal, use?	Shared presentations?	Trauma, sleep, grief, stress?

Stop trying to prove your first impression — start trying to challenge it.

6 Confidence isn't never changing your mind

INITIAL FORMULATION → NEW INFORMATION → UPDATED FORMULATION

What can change the picture: new history, collateral from family or records, new symptoms, and treatment response.



Changing the diagnosis ≠ getting it wrong. The diagnosis should follow the evidence — not your ego.

7 Confidence ≠ certainty

Confidence can sound like:

"This is my leading diagnosis." *"I need more history."* *"I want collateral."*

"These are the differentials I'm still considering."

"I need to rule something else out." *"I'm going to reassess this over time."*

FALSE CERTAINTY
"I know exactly what this is."

≠

CLINICAL CONFIDENCE
"I can explain what I think, why, and what could change my mind."

8 The goal isn't to diagnose faster

Speed and confidence are not the same thing. Learn when you have enough information:

ASK	Have I established a clear timeline?
COMPARE	Have I considered reasonable differentials?
RULE OUT	Medical, medication, and substance causes?
CONTEXTUALIZE	Do symptoms fit history and functioning?
REASSESS	Do I need more time or information?

Sometimes slowing down IS the clinical skill. Not every diagnosis needs to be finalized in one visit.



How diagnostic confidence is built

Repetition · Reflection · Mistakes · Supervision · Continued learning · Seeing patients over time

Give yourself time.

"This is what I think based on what I know today — and I'm willing to reassess when I know more."

Thoughtful without every answer. Confident without certainty. Decisive without rigidity.

ADAPTED FROM @PSYCHIATRYWITHKIARA