Understanding the

Temporomandibular Joints for Proper Assessment *and* Treatment Planning

James R. McKee, DDS

Abstract

Many patients who are seeking esthetic enhancements also present with occlusal issues. As a profession, we have assumed occlusal issues are due to improper tooth position, as this causes occlusal disharmonies resulting in problems that can present as wear, mobility, or sensitivity at the tooth level. Three-dimensional imaging offers new insights into many of the occlusal problems patients present with today. Many of the tooth position problems we see clinically are the result of structural changes in the temporomandibular (TM) joints. This article outlines some key points regarding occlusion and the TM joints.

Key Words: TM joints, magnetic resonance imaging, cone beam computed tomography, condylar position, occlusion

Bonus content! This article has a CE course on AACD's Virtual Campus. To view, go to AACD.com/virtualcampus While these diagnostic modalities offer valuable information, they have one common trait: They all are based on an indirect visualization of TM joint anatomy.

Kin hin

66

Introduction

It is always exciting when patients seek treatment to improve the esthetic appearance of their teeth. The excitement, however, often turns to uncertainty—and, in some cases, anxiety—when the patient also reports having "clicking" jaw joints. Several questions arise in this scenario: Are the clicking joints a problem? Do they need to be treated? *Can* they be treated? Is magnetic resonance imaging (MRI) or cone beam computed tomography (CBCT) necessary? Is it possible to fabricate an occlusal appliance to treat the clicking joints?

Assessing the Joints

The uncertainty about clicking jaw joints is related to how we have been taught to assess temporomandibular (TM) joint condition. The common clinical exam to assess the joints includes taking a history; measuring range of motion; palpating muscles; and load testing and listening to the joints using a stethoscope, doppler auscultation, or joint vibration analysis. While these diagnostic modalities offer valuable information, they have one common trait: They all are based on an indirect visualization of TM joint anatomy (Fig 1). As a result, there is uncertainty about what to tell patients regarding both treatment options and the prognosis of these treatment options, because we do not have a clear understanding of the anatomy we are treating.

Imaging

Imaging provides direct visualization of the TM joints. The ability to directly assess the soft tissue (disc) with MRI¹⁻⁷ and hard tissue (condyle/joint socket) with CBCT⁸⁻¹⁰ offers both dentists and patients a clear understanding of joint anatomy. This leads to a more realistic discussion about treatment options and the prognosis for successful treatment.

MRI: The TM joint MRI begins with assessing disc position. Historically, normal disc position was determined as $12:00 \pm 10^{0.11}$ from the superior surface of the condyle (Fig 2). However, the definition of "normal" changed in 1997 from $12:00 \pm 100$ to $12:00 \pm 30^{0.12}$ (Fig 3). In reality, disc position is best evaluated based on the disc's ability to dissipate the load that is applied to the TM joints. The optional load-bearing position of the disc is 11:00, with normal anterior and posterior aspects of the disc on each side of the 11:00 load-bearing position.¹³ With this optimal anatomy for load bearing, the posterior attachment for a structurally intact or normal TM joint is at approximately the 1:00 position, as opposed to the assumed normal 12:00 position (Fig 4).



Figure 1: TM joint diagram.





Figure 2: TM joint MRI (12:00 ± 10°).

Figure 3: TM joint MRI (12:00 ± 30°).



Loading bearing position 11:00 Posterior attachment 1:00

Figure 4: TM joint MRI (11:00 load bearing/1:00 posterior attachment).



Figure 5: MRI, coronal view.



Figure 6: CBCT, sagittal view.



Figure 7: CBCT, coronal view.



Figure 8: CBCT, sagittal view.



Figure 9: CBCT showing intact and non-intact cortical plate.



Figure 10: Normal ramus length.

The disc is designed to be compressed during function. As a result, it receives its nutritional components through synovial fluid. In order for the disc to receive nutrition, it must be in the correct position. Disc position is dictated by the ligament attachment of the disc to the condyle (Fig 5). The disc attaches at the condyle's lateral, medial, and posterior aspects. Normal disc anatomy serves to correctly position the condyle in the joint socket. In the adult patient, normal disc anatomy also maintains the vertical dimension of the TM joint, which maintains the occlusion. In growing patients, normal disc position fosters growth to the full genetic potential.

CBCT: CBCT imaging enables assessment of the hard tissue in the TM joint. Normal condyle size is approximately 8 mm in an anterior-posterior dimension (Fig 6) and 20 mm in a medial-lateral perspective (Fig 7). CBCT imaging also allows for an appreciation of the heavily buttressed bone at the medial aspect of the joint socket. This bone provides ideal anatomy in the joint socket for load distribution from the masticatory muscles. In addition, CBCT imaging facilitates evaluation of condylar position in the joint socket (Fig 8) as well as the ability to analyze the integrity of the condylar cortical plate (Fig 9). Lastly, CBCT imaging allows for an evaluation of the ramus length. Normal ramus length is approximately 60 to 70 mm (Fig 10).

TM Joint Structural Changes

Ligament Injuries

Structural changes in the TM joints begin with an injury to the soft tissue attachment of the disc to the condyle. The ligament injury typically is caused by either a compression, a stretch, or a whiplash injury to the TM joint.14,15 The ligament damage can result in a clicking joint if the disc maintains its shape despite not being in the correct position to receive synovial fluid. This type of injured joint is referred to as a clicking joint or a disc displacement with reduction (Fig 11). Clicking joints can stop clicking when the disc loses its natural biconcave shape due to a lack of synovial fluid compression. This type of injured joint is referred to as a locking joint or a disc displacement without reduction (Fig 12). In both scenarios, the condyle compresses the retrodiscal tissue as opposed to disc tissue. The advanced stage of joint breakdown occurs when the condyle perforates the retrodiscal tissue and there is direct bone-to-bone contact between the condyle and the joint socket (Fig 13).

In addition to assessing the ligament attachment from a sagittal perspective, it is important to assess it from a coronal perspective. Some patients will present with an injury to the lateral pole ligament attachment while maintaining the medial pole attachment. Injuries at the lateral pole can result in a clicking joint (lateral pole disc displacement with reduction) or a locking joint (lateral pole without reduction). Other patients will present with a lateral and medial pole displacement with or without reduction.¹⁶⁻²²

Risk Levels

From a clinical perspective, normal joints present with the least amount of risk for both the patient and the dentist. Patients with normal TM joints have normal hard and soft tissue. The likelihood for significant pain issues or bite instability is low. Patients also can present with joint injuries confined to the lateral pole. These patients tend to have relatively low risk factors since the medial pole of the condyle is still covered by normal disc tissue.

Pain: The highest level of risk occurs when the ligament attachment is injured at the lateral and the medial pole. Typically, TM joint changes are defined through pain. While pain is usually assumed to come from masticatory muscles, TM joint imaging has provided insight regarding different sources of pain. Given the insights from three-dimensional (3D) imaging, it is clear pain can come from herniated discs (Fig 14) impinging on different anatomic structures. Condylar changes such as eroded bone (Fig 15), small bone (Fig 16), or edematous bone (Fig 17) also can cause significant pain.²³⁻³² Pain can occur from structural changes in the upper cervical spine as well as from elevated sympathetic nerve conduction.³³⁻³⁶

Occlusion: Structural changes in the TM joint can cause not only pain, but also significant changes in the occlusion. The loss of vertical dimension from structural changes in the TM joint can present several considerable issues for the patient seeking cosmetic improvements from an occlusal perspective.



Figure 11: Disc displacement with reduction.









Figure 13: Disc displacement with perforation.



Figure 15: Eroded condylar bone.



Figure 14: Herniated TM joint disc.

Figure 16: Small condylar bone.

Figure 17: Edematous condylar bone.



Figure 18: Class II occlusion with short rami.



Figures 19a & 19b: Anterior tooth uncoupling.

These occlusal changes usually present as a Class II bite shift due to a loss of soft tissue dimension at the disc level and a loss of hard tissue dimension at the condyle level. The greater the loss of vertical dimension, the greater the likelihood of a Class II bite shift (Fig 18). Loss of vertical dimension at the TM joint level can occur though joint degeneration of a normally developed TM joint. The more common loss of vertical dimension at the joint level occurs through a lack of complete growth due to joint injuries in the growing patient.³⁷⁻⁴⁸ If the disc ligaments are compromised in the growing patient, the protective nature of the disc is lost and both mandibular and maxillary growth are compromised. The typical clinical presentation is the uncoupling of the anterior teeth (Figs 19a & 19b).

The key to understanding the role occlusion plays in both pain and bite issues is 3D imaging of the TM joints' soft and hard tissue. The MRI can provide diagnostic information related to disc position and disc condition as well as evaluate the condition of the marrow space to assess condylar edema. The CBCT can provide diagnostic information related to condylar size and cortical plate integrity. It also can offer diagnostic information related to ramus length, airway, and upper cervical spine issues.

The MRI is obtained using a multipositional condylar protocol developed by Dr. Mark Piper **(Table 1)**. The condylar positions are fully seated in maximum intercuspation, at incisal edge position, and at translated condylar position of 30 mm. The purpose of imaging in different condylar positions is to

66

In reality, disc position is best evaluated based on the disc's ability to dissipate the load that is applied to the TM joints.

Table 1. Piper MRI Protocol

PIPER CLINIC TMJ MRI SCAN PROTOCOL FULL

Orientation	Sequence	Occlusion	Starting Side	View	Starting Point	Ending Point	Qty	Trs	Hfs Sp
Axial	Tı Localizer	None	Bilateral	Inf To Sup	Mandibular Ramus	Superior Orbit	16	240/4.0	5.0
Axial	T2	None	Bilateral	Sup To Inf	Superior Orbit	Mandibular Ramus	20	180/3.0	3.0
Coronal	T2 Localizer	None	Bilateral	Post To Ant	Eac Anterior	Orbit Posterior	14	200/4.0	4.0
Sagittal	Tı	Bite 1 Fscp	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Coronal	Tı	Bite 1 Fscp	Left Then Right	Post To Ant	Eac Anterior	Anterior To Eminence	18	140/3.0	2.8
Sagittal	Stir	Bite 1 Fscp	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Coronal	T2	Bite 1 Fscp	Left Then Right	Post To Ant	Eac Anterior	Anterior To Eminence	18	140/3.0	2.8
Sagittal	Proton Density	Bite 2 lep	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Coronal	Proton Density	Bite 2 lep	Left Then Right	Post To Ant	Eac Anterior	Anterior To Eminence	18	140/3.0	2.8
Sagittal	Proton Density	Prop Open Tcp	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Coronal	Proton Density	Prop Open Tcp	Left Then Right	Post To Ant	Eac Anterior	Anterior To Eminence	18	140/3.0	2.8
Sagittal	Proton Density	Closed Mouth Mip	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8

PIPER CLINIC TMJ MRI SCAN PROTOCOL BASIC

Orientation	Sequence	Occlusion	Starting Side	View	Starting Point	Ending Point	Qty	Trs	Hfs Sp
Axial	T2	None	Bilateral	Sup To Inf	Superior Orbit	Mandibular Ramus	20	180/3.0	3.0
Sagittal	Тı	Bite 1 Fscp	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Sagittal	Stir	Bite 1 Fscp	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Coronal	T2	Bite 1 Fscp	Left Then Right	Post To Ant	Eac Anterior	Anterior To Eminence	18	140/3.0	2.8
Sagittal	Proton Density	Bite 2 lep	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Sagittal	Proton Density	Prop Open Tcp	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8
Sagittal	Proton Density	Closed Mouth Mip	Left Then Right	Lat To Med	Lateral To Condyle	Medial To Condyle	16	140/3.0	2.8

PIPER CLINIC TMJ MRI SCAN PROTOCOL SCREEN

Orientation	Sequence	Occlusion	Starting Side	View	Starting Point	Ending Point	Qty	Trs	Hfs Sp
Axial	T2	None	Bilateral	Sup To Inf	Superior Orbit	Mandibular Ramus	20	180/3.0	3.0
Sagittal	Stir	Bite 1 Fscp	Left Then Right	Lat To Med	Lateral Condyle	Medial To Condyle	16	140/3.0	2.8
Coronal	T2	Bite 1 Fscp	Left Then Right	Post To Ant	Eac Anterior	Anterior Eminence	18	140/3.0	2.8
Sagittal	Proton Density	Bite 2 lep	Left Then Right	Lat To Med	Lateral Condyle	Medial To Condyle	16	140/3.0	2.8



Figures 20a-20c: Piper MRI condylar positioning indices.

assess the positions' condyle–disc interface. The images in different condylar position are obtained using occlusal indices to position the mandible during the MRI (Figs 20a-20c). The CBCT is obtained in a fully seated condylar position, as are the diagnostic study casts and digital photographs. Obtaining diagnostic records in the same condylar position allows for cross-referencing the different information that can be gained from the different diagnostic modalities.

Summary

Understanding TM joint condition allows for an increased level of confidence when discussing potential cosmetic changes with patients. The dentist's ability to assess the risk from structural changes in the soft and hard tissue provides invaluable data when developing a treatment plan and enables the dentist to help patients make treatment decisions based on accurate, precise information. Gaining knowledge of the anatomy through 3D imaging can help dentists have confidence when patients ask about their "clicking" jaw joints.

References

- Schellhas KP, Wilkes CH, Fritts HM, Omlie MR, Heithoff KB, Jahn JA. Temporomandibular joint: MR imaging of internal derangements and postoperative changes. Am J Roentgenol. 1988 Feb;150(2):381-9.
- Schellhas KP, Wilkes CH. Temporomandibular joint inflammation: comparison of MR fast scanning with T1- and T2-weighted imaging techniques. Am J Roentgenol. 1989 Jul;153(1):93-8.
- Lieberman JM, Hans MG, Rozencweig G, Goldberg JS, Bellon EM. MR imaging of the juvenile temporomandibular joint: preliminary report. Radiology 1992 Feb;182(2):531-4.
- Larheim TA. Role of magnetic resonance imaging in the clinical diagnosis of the temporomandibular joint. Cells Tissues Organs. 2005;180(1):6-21.
- Whyte AM, McNamara D, Rosenberg I, Whyte AM. Magnetic resonance imaging in the evaluation of temporomandibular joint disc displacement a review of 144 cases. Int J Oral Maxillofac Surg. 2006 Aug;35(8):696-703.
- Tomas X, Pomes J, Berenguer J, Quinto L, Nicolau C, Mercader JM, Castro V. MR imaging of temporomandibular joint dysfunction: a pictorial review. Radiographics. 2006 May-Jun;26(3):765-81.
- Matsubara R, Yanagi Y, Oki K, Hisatomi M, Santos KC, Bamgbose BO, Fujita M, Okada S, Minagi S, Asaumi J. Assessment of MRI findings and clinical symptoms in patients with temporomandibular joint disorders. Dentomaxillofac Radiol. 2018 May;47(4):20170412.
- Larheim TA. Current trends in temporomandibular joint imaging. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 1995 Nov;80(5):555-76.
- Alomar X, Medrano J, Cabratosa J, Clavero JA, Lorente M, Serra I, Monill JM, Salvador A. Anatomy of the temporomandibular joint. Semin Ultrasound, CT MR. 2007 Jun;28(3):170-83.
- Larheim TA, Abrahamsson AK, Kristensen M, Arvidsson LZ. Temporomandibular joint diagnostics using CBCT. Dentomaxillofac Radiol. 2015;44(1):20140235.
- Rammelsberg P, Pospiech PR, Jager L, Pho Duc JM, Bohm AO, Gernet W. Variability of disc position in asymptomatic volunteers and patients with internal derangements of the TMJ. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 1997 Mar;83(3):393-9.
- Drace JE, Enzmann DR. Defining the normal temporomandibular joint: closed-, partially open-, and open-mouth MR imaging of asymptomatic subjects. Radiology. 1990 Oct; 177(1): 67-71.
- Chang MS, Choi JH, Yang IH, An JS, Heo MS, Ahn SJ. Relationships between temporomandibular joint disc displacements and condylar volume. Oral Surg Oral Med Oral Pathol Oral Radiol. 2018 Feb;125(2):192-8.

- 14. Harkins SJ, Marteney JL. Extrinsic trauma: a significant precipitating factor in temporomandibular dysfunction. J Prosthet Dent. 1985 Aug;54(2):271-2.
- 15. Schellhas P. Temporomandibular joint injuries. Radiology. 1989 Oct;173(1):211-6.
- Katzberg RW, Tallents RH, Hayakawa K, Miller TL, Goske MJ, Wood BP. Internal derangements of the temporomandibular joint: findings in the pediatric age group. Radiology. 1985 Jan;154(1):125-7.
- Wilkes CH. Internal derangements of the temporomandibular joint. Pathological variations. Northwest Dent. 1990 Mar-Apr;69(2)25-32.
- Paesani D, Westesson PL, Hatala M, Tallents RH, Kurita K. Prevalence of temporomandibular joint internal derangement in patients with craniomandibular disorders. Am J Orthod Dentofacial Orthop. 1992 Jan;101(1):41-7.
- Nebbe B, Major PW. Prevalence of TMJ disc displacement in a pre-orthodontic adolescent sample. Angle Orthod. 2000 Dec;70(6):454-63.
- Tomas X, Pomes J, Berenguer J, Mercader JM, Pons F, Donoso L. Temporomandibular joint soft-tissue pathology, II: nondisc abnormalities. Semin Ultrasound CT MR. 2007 Jun;28(3):205-12.
- Molinari F, Manicone PF, Raffaelli L, Raffaelli R, Pirronti T, Bonomo L. Temporomandibular joint soft-tissue pathology, I: disc abnormalities. Semin Ultrasound CT MR. 2007 Jun;28(3):192-204.
- Ikeda K, Kawamura A. Disc displacement and changes in condylar position. Dentomaxillofac Radiol. 2013 Mar;42(3):84227642.
- Westesson PL. Structural hard-tissue changes in temporomandibular joints with internal derangement. Oral Surg Oral Med Oral Pathol. 1985 Feb;59(2):220-4.
- Kurita H, Ohtsuka A, Kobayashi H, Kurashina K. Alteration of the horizontal mandibular condyle size associated with temporomandibular joint internal derangement in adult females. Dentomaxillofac Radiol. 2002 Nov;31(6):373-8.
- 25. Kurita H, Kojima Y, Nakatsuka A, Koike T, Kobayashi H, Kurashina K. Relationship between temporomandibular joint (TMJ)-related pain and morphological changes of the TMJ condyle in patients with temporomandibular disorders. Dentomaxillofac Radiol. 2004 Sep;33(5):329-33.
- 26. Sano T, Otonari-Yamamoto M, Otonari T, Yajima A. Osseous abnormalities related to the temporomandibular joint. Semin Ultrasound CT MR. 2007 Jun;28(3):213-21.
- Kung JW, Yablon CM, Eisenberg RL. Bone marrow signal alteration in the extremities. Am J Roentgenol. 2011 May;196(5):492-510.

- 28. Hasegawa H, Saitoh I, Nakakura-Ohshima K, Shigeta K, Yoshihara T, Suenaga S, Inada E, Iwasaki T, Matsumoto Y, Yamasaki Y. Condylar shape in relation to anterior disc displacement in juvenile females. Cranio. 2011 Apr;29(2):100-10.
- Larheim TA, Sano T, Yotsui Y. Clinical significance of changes in the bone marrow and intra-articular soft tissues of the temporomandibular joint. Semin Orthod. 2012 Mar;18(1):30-43.
- 30. Dias IM, Coelho PR, Assis NMSP, Leite FPP, Devito KL. Evaluation of the correlation between disc displacements and degenerative bone changes of the temporomandibular joint by means of magnetic resonance images. Int J Oral Maxillofac Surg. 2012 Sep;41(9):1051-7.
- Talaat W, Al Bayatti S, Al Kawas S. CBCT analysis of bony changes associated with temporomandibular disorders. Cranio. 2016 Mar;34(2)88-94.
- Bae S, Park MS, Han JW, Kim YJ. Correlation between pain and degenerative bony changes on cone-beam computed tomography images of temporomandibular joints. Maxillofac Plast Reconstr Surg. 2017 Jul;39(1):19.
- Fechir M, Geber C, Birklein F. Evolving understandings about complex regional pain syndrome and its treatment. Curr Pain Headache Rep. 2008 Jun;12(3):186-91.
- 34. Harden RN, Oaklander AL, Burton AW, Perez RS, Richardson K, Swan M, Barthel J, Costa B, Graciosa JR, Bruehl S. Complex regional pain syndrome: practical diagnostic and treatment guidelines, 4th edition. Pain Med. 2013 Feb;14(2):180-229.
- Tinastepe N, Oral K. Complex regional pain syndrome. J Am Dent Assoc. 2015 Mar;146(3):200-2.
- Parkitny L, Wand BM, Graham C, Quintner J, Moseley GL. Interdisciplinary management of complex regional pain syndrome of the face. Phys Ther. 2015 Jul;96(7):1067-73.
- 37. Schellhas KP. Unstable occlusion and temporomandibular joint disease. J Clin Orthod. 1989 May;23(5):332-7.
- Schellhas P, Keck RJ. Disorders of skeletal occlusion and temporomandibular joint disease. Northwest Dent. 1989 Jan-Feb;68(1):35-9, 42.
- Schellhas KP, Piper MA, Omlie MR. Facial skeleton remodeling due to temporomandibular joint degeneration: an imaging study of 100 patients. Am J Roentgenol. 1990 Aug;155(2):373-83.



- 40. Nebbe B, Major PW, Prasad NG. Female adolescent facial pattern associated with TMJ disc displacement and reduction in disc length: part I. Am J Orthod Dentofacial Orthop. 1999 Aug;116(2):168-76.
- 41. Qadan S, Macher DJ, Tallents RH, Kyrkanides S, Moss ME. The effect of surgically induced anterior disc displacement of the temporomandibular joint on the midface and cranial base. Clin Orthod Res. 1999 Aug;2(3):124-32.
- 42. Paesani D, Salas E, Martinez A, Isberg A. Prevalence of temporomandibular joint disc displacement in infants and young children. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 1999 Jan;87(1):15-9.
- Defabianis P. Post-traumatic TMJ internal derangement: impact on facial growth (findings in a pediatric age group). J Clin Pediatr Dent. 2003 Summer;27(4):297-303.
- 44. Flores-Mir C, Nebbe B, Heo G, Major PW. Longitudinal study of temporomandibular joint disc status and craniofacial growth. Am J Orthod Dentofacial Orthop. 2006 Sep;130(3):324-30.
- 45. Emshoff R, Moriggl A, Rudisch A, Laimer K, Neunteufel N, Crismani A. Are temporomandibular joint disc displacements without reduction and osteoarthrosis important determinants of mandibular backward positioning and clockwise rotation? Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2011 Apr;111(4):435-41.
- 46. Maglione H, de Zavaleta LA, Laraudo J, Falisi G, Fernandez F. Temporomandibular dysfunction: internal derangement associated with facial and/or mandibular asymmetry. Cranio. 2013 Oct;31(4):276-82.
- 47. Manfredini D, Segu M, Arveda N, Lombardo L, Siciliani G, Rossi A, Guarda-Nardini L. Temporomandibular joint disorders in patients with different facial morphology. A systematic review of the literature. J Oral Maxillofac Surg. 2016 Jan;74(1):29-46.
- 48. You KH, Kim KH, Lee KJ, Baik HS. Three-dimensional computed tomography analysis of mandibular morphology in patients with facial asymmetry and mandibular retrognathism. Am J Orthod Dentofac Orthop. 2018 May;153(5):685-91. **jCD**



Dr. McKee is in private practice in Downers Grove, Illinois. He also is a visiting faculty member at Piper Education and Research Center and a resident faculty member at Spear Education.

Disclosure: The author did not report any disclosures.

