


A detectability criterion for sequential data assimilation

Jason Frank

Mathematical Institute
Utrecht University

SIAM, Snowbird
May 23, 2019



A detectability criterion for sequential data assimilation

Jason Frank

Mathematical Institute
Utrecht University



with
Sergiy Zhuk
IBM Dublin

SIAM, Snowbird
May 23, 2019

Motivation

Originally motivated to understand what minimal observations are needed to construct a complete state estimation.

We construct a continuous time synchronous data assimilation method that explicitly uses a decomposition of the tangent space into expanding and decaying subspaces.

Lyapunov vectors and data assimilation in the literature:

- AUS methods (Trevisan, Carrassi, Bocquet, Grudzien, ...)
- Mentioned in work by Ghil et al., González Tokman & Hunt, Gottwald & Reich, ...
- *De Leeuw, Dubinkina, Frank, Steyer, Tu, Van Vleck 2018.*

Synchronization of chaotic dynamics

Driver:

Observation time series:

$$\text{Driver:} \quad \dot{\tilde{x}} = f(t, \tilde{x}) \quad y(t) = H\tilde{x}(t) \quad (t)$$

$$\text{Receiver:} \quad \dot{x} = f(t, x(t)) + \alpha L(t)(y(t) - Hx(t))$$

Transverse linear dynamics:

$$w(t) = x(t) - \tilde{x}(t)$$

$$\dot{w} = [A(t) - \alpha L(t)H] w + N(t, w)$$

$$A(t) = Df(x(t))$$

$$\text{Sync:} \quad \lim_{t \rightarrow \infty} \|w(t)\| \leq Ce^{-\mu t}$$



$$\lim_{t \rightarrow \infty} \|x(t) - \tilde{x}(t)\| \leq Ce^{-\mu t}$$

(e.g. Pecora & Carroll 1990)

Lyapunov exponents and stability

(1) Linear systems

$x(t)=0$ is exponentially stable if

$$\dot{x}(t) = A(t)x(t)$$

$$\lambda_1 < 0$$

2. Quasi-linear systems

$x(t)=0$ is stable in a
neighborhood of 0 if

(1) holds

$$\dot{x} = A(t)x + N(t, x), \quad N(t, 0) = 0$$

$$\|N(t, x) - N(t, \tilde{x})\| \leq C\|x - \tilde{x}\|^p, \quad p > 1$$

3. Nonlinear systems

Orbit $x^*(t)$ attracts a
neighborhood of itself
if (1) holds

$$\dot{x} = f(t, x), \quad x(t) = x^*(t) + w(t)$$

$$\begin{aligned} \dot{w} &= Df(x^*)w + [f(x) - f(x^*) - Df(x^*)w] \\ &= A(t)w + N(t, w) \end{aligned}$$

Computation of Lyapunov Exponents

Procedure to compute the LEs yields an orthogonal basis for the associated Lyapunov vectors.

Continuous QR factorization $\dot{X}(t) = A(t)X(t), \quad X(t) = Q(t)R(t)$

$$\dot{R} = BR$$

$$B = Q^T A Q - S$$

$$S = -S^T$$

$$\dot{Q} = (I - QQ^T)AQ - QS$$

$$B = \begin{bmatrix} B_{11} & * & * \\ & \ddots & * \\ & & B_{kk} \end{bmatrix}$$

$$S = \begin{bmatrix} 0 & -\text{tril}(Q^T A Q) \\ \text{tril}(Q^T A Q) & 0 \end{bmatrix}$$

In essence a power iteration.

First k columns of Q span the k fastest growing Lyapunov vectors. Q is $d \times k$.

LEs appear ordered on the diag of B :

$$\lambda_i = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t B_{ii}(s) ds$$

Sequential DA with tangent splitting

Receiver / filter: $\dot{x} = f(t, x(t)) + \alpha L(t)(y(t) - Hx(t))$

Transverse dynamics:

$$w(t) = x(t) - \tilde{x}(t)$$
$$\dot{w} = [A(t) - \alpha L(t)H] w + N(t, w)$$
$$A(t) = Df(x(t))$$

Note that $\tilde{x}(t)$ solves the filter equation.

Choose $L(t)$ such that the linearization has $\lambda_1 < 0$, then the filter exponentially attracts a neighborhood of itself.

$$B = Q^T A Q - S$$

$$\begin{aligned}\tilde{B} &= Q^T (A - \alpha L H) Q - S \\ &= B - \alpha Q^T L H Q \\ &= B - \alpha \tilde{R}\end{aligned}$$

Choose $\tilde{R} = Q^T L H Q$ upper triangular with positive diagonal!

Sequential DA with tangent splitting

Receive

Want $\tilde{R} = Q^T L H Q$ upper tri. positive diag.

Transv

Choose $L = Q U H^T$
then $\tilde{R} = Q^T Q U H^T H Q$
 $= U H^T H Q$

, w)

Suppose U is invertible: $U^{-1} \tilde{R} = H^T H Q$

A **good choice** is the QR factorization:

$$\tilde{Q} \tilde{R} = H^T H Q$$

Consequently, $U = \tilde{Q}^T$

$$L = p Q \tilde{Q}^T H^T$$

Note that
Choose
then the
itself.

$$B = Q$$

upper

triangular with positive diagonal!

Detectability criterion

Recall $\tilde{Q}\tilde{R} = H^T H Q$

The Lyapunov vector v_j is *detectable* if

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \int_0^t \tilde{R}_{ii}(s) ds > 0, \quad i = 1, \dots, j$$

The *pair* $(A(t), H)$ is detectable if all Lyapunov vectors corresponding to nonnegative Lyapunov exponents are detectable. (*Needs rank $H \geq$ dimension nonstable space*).

Theorem. If $(A(t), H)$ is detectable then there exists $\alpha > 0$ such that all Lyapunov exponents of the fundamental matrix equation

$$\dot{W} = (A(t) - \alpha L(t)H)W, \quad L(t) = Q(t)\tilde{Q}(t)^T H^T$$

are negative.

Observation operator

By construction H^T and Q have full rank. Because they are 'tall' matrices, they have no null space.

Consequently, $H^T H Q x = 0$ implies $H Q x = 0$. But $H Q$ has dim. $m \times k$ and rank $\min\{m, k\}$ unless there is a nontrivial linear combination of the columns of Q in $\ker(H)$, in which case H doesn't 'see' the whole nonstable space.


Conclusion: We need $m \geq k$ and $H Q$ full rank for detectability.

Numerical experiment

- Lorenz 96 model, $M=18$ lattice points, initial condition

$$x_j(0) = \sin 2\pi j/M + \mathcal{N}(0, \epsilon^2)$$

100-member ensemble, $H =$ first 8 eigenmodes



Numerical experiment

- Lorenz 96 model, $M=18$ lattice points, initial condition
- Noisy data: $\eta(t) \sim \mathcal{N}(0, 4)$




Figure 4: Comparison of the filter (29) and the ExKF (33) for the Lorenz '96 model (32) with $k = 8$. Left, the errors $\|\xi(t)\|$ for a 10-member ensemble of perturbed initial conditions. Right, the errors $\|\xi(t)\|$ for a 10-member ensemble with random observational error.